



Oilseed Meals an Untapped Resource for Nutritional Innovation

Karnika ^{a,b++*} and Asha Kawatra ^a

^a Department of Foods and Nutrition, Chaudhary Charan Singh Haryana Agricultural University, Hisar, 125001, India.

^b Punjab Agricultural University, Ludhiana, 141004, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i122734>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129108>

Original Research Article

Received: 26/10/2024

Accepted: 30/12/2024

Published: 30/12/2024

ABSTRACT

The present investigation was carried out to nutritionally evaluate *laddo* supplemented with oilseed meals, including sesame meal (SM), flaxseed meal (FM), and groundnut meal (GM) at three level of incorporation (12, 18, and 24%). Composite meal was also prepared by mixing sesame, flaxseed & groundnut meal in equal amounts for the development of *laddo* similarly at three levels of incorporation (12, 18, and 24%). All the developed *laddo* were found to be organoleptically acceptable adjudicated by nine-point hedonic scale. Acceptability scores of all kinds of *laddo* varied between 'liked moderately' to 'liked very much' category. *Laddo* with 24% of incorporation of oilseed meal were studied for nutritional composition. Corresponding *laddo* prepared without oilseed meal was taken as control. Incorporation of all the three oilseed meals individually and in combination resulted in significant improvement in level of crude protein, fat, ash, and crude fiber, which ranged from 13.35-20.03, 25.52-35.99, 3.97-5.70, 4.19-5.14%, respectively. One serving (25gm) of *laddo* can meet 27% requirement of protein for pre-schoolers as per RDA (Recommended Dietary

⁺⁺Presently Pursuing Ph.D. in Food and Nutrition;

*Corresponding author: E-mail: chaudhary.karnika12@gmail.com;

Allowances) 2020. Value added *Laddo* if promoted and popularized will help in the utilization of oilseed meals which are generally discarded and have a great nutritive potential which can be used for enrichment of daily diet at a reasonable cost. The developed recipe can be efficiently used as a means of improving nutritional status and to eliminate the malnutrition.

Keywords: Oilseed meals; laddo and malnutrition.

1. INTRODUCTION

After the extraction of oil from oilseeds, the residue (by-products of oil extraction) is obtained. Those residues are known as oilseed meal/oilseed cake which are used as fodder or are discarded as waste. From one ton of oilseed approx. 500 to 650 kilos of the meal is obtained depending on the type of oilseed and as by practice half of the oilseed cake received from the production is being dissipated. Oilseed meals are nutritionally rich and major sources of protein, fatty acids, polyphenols, and dietary fibre. There are several oilseed meals available like rape & mustard seed meal, sesame seed meal, flaxseed meal, groundnut seed meal, etc. All these are nutrient-dense and very economical and can be used to develop human foods that can be directly consumed (Karnika et al. 2022).

Peanut also known as groundnut (*Arachis hypogaea*) is a legume crop grown mainly for its edible seeds. Due to its high oil content, it is classified as a grain and legume and an oil crop (Dwiwedi et al. 2011). Peanut flour obtained after dehulling and grinding contained 49.97% crude protein, 2.80% crude fiber, 3.50% ash and 27.54% fat (Park et al. 2017).

Flaxseed (*Linum usitatissimum*) also known as linseed, is a flowering plant in the family of Linaceae. After the oil extraction flaxseed meal is obtained which is protein-rich and can be used in different ways for human use. This meal is rich in vitamin B₆ and contains fiber up to 2-7%. Vitamin B₆ is an essential nutrient which body uses as a coenzyme in some 100s of enzymatic reactions in the human body (Heuze et al. 2017).

Sesame (*Sesamum indicum*) is a flowering plant in the genus *Sesamum*, also known as *benne* (Merriam-Webster.com 2021). Sesame oil meal is a protein-rich by-product. Expeller sesame meal has a protein content of about 45% and ranging from 32 to 53% whereas solvent extracted sesame meal contains about 48% protein (Karnika et al. 2022).

Considering the nutritional potential of selected oilseed meals which are otherwise discarded as

waste, present investigation was carried out to develop nutrient dense *laddo* which can be made easily and popularized among all age groups. The value added *laddo* can be used for enrichment of daily diet which can be helpful for improving nutritional status of community and may contribute to eradicate problem of malnutrition.

2. MATERIALS AND METHODS

The present study was carried out in the Department of Foods and Nutrition, I.C. College of Home Science, Chaudhary Charan Singh Haryana Agricultural University, Hisar. Sesame and peanut meal were procured from the local market, and flaxseed meal (cold-pressed) was procured from India Mart.

The required ingredients for the preparation of *Laddo* namely, whole wheat flour, ghee, refined sugar, peanut & sesame were purchased in a single lot from the local market of Hisar.

2.1 Product Development

A spherical treat from the Indian subcontinent, *laddu* or *laddo* is created with a variety of ingredients and either sugar syrup or jaggery. According to some, it's "possibly the most ancient and universal of Indian sweets (*The Bloomsbury Handbook of Indian Cuisine* 2023).

Twelve types of *laddo* were prepared by replacing whole wheat flour at 12 % (T1 & C1), 18 % (T2 & C2) and 24 % (T3 & C3) of wheat flour by sesame meal, peanut meal, flaxseed meal and composite meal (prepared by mixing sesame, flaxseed, and groundnut meal in equal amount) individually (Table 1). Corresponding *laddo* prepared without the addition of oilseed meals were served as control. For the control *laddo*, standard recipe ingredient proportions of wheat flour, ground sugar, ghee, roasted sesame seeds, and roasted peanut were 50 g, 30 g, 30 g, 15 g, and 25 g, respectively. Wheat flour and oilseed meal powder was sieved separately and oilseed meal was dry roasted whereas wheat flour was roasted in ghee. After roasting of wheat

Table 1. Types of laddo

Incorporated recipe	Types of laddo	Level of incorporation
Control	control	Whole wheat flour::100
Sesame meal	T1	Whole wheat flour:SM::88:12
	T2	Whole wheat flour:SM::82:18
	T3	Whole wheat flour:SM::76:24
Flaxseed meal	T1	Whole wheat flour:FM::88:12
	T2	Whole wheat flour:FM::82:18
	T3	Whole wheat flour:FM::76:24
Groundnut meal	T1	Whole wheat flour:GM::88:12
	T2	Whole wheat flour:GM::82:18
	T3	Whole wheat flour:GM::76:24
Composite meal	C1	Whole wheat flour:CM::88:12
	C2	Whole wheat flour:CM::82:18
	C3	Whole wheat flour:CM::76:24

Control- without any oilseed meal incorporation (T1 & C1, T2 & C2 and T3 & C3 are 12%, 18%, and 24%)

Flour and oilseed meals, roasted and crushed peanut and sesame were added to the mixture and removed from fire. Ground sugar was added after mixture was slightly cooled down. Balls were prepared from the mixture.

2.2 Organoleptic Evaluation

Organoleptic evaluation of supplemented oilseed meal *Laddo* adjudicated using nine point hedonic rate scale by a panel of ten semi-trained judges from I.C College of Home Science, CCS Haryana Agricultural University, Hisar. The judges were presented with each oilseed meal *laddo* at separate time session along with control. Organoleptic evaluation were made with respect to color, appearance, aroma, texture, and taste on a scale of one to nine representing 'dislike extremely' to 'like extremely' categories. The scale of five represented the neutral score 'neither like nor dislike'.

2.3 Nutritional Evaluation

Moisture, ash, crude fat, crude protein, and crude fiber were estimated using the standard analysis methods (AOAC, 2010).

2.4 Statistical Analysis

The data obtained from nutritional analysis and organoleptic evaluation was statistically analyzed using mean, standard error, and ANOVA (one-way and two-way analysis). Replicates were used for experiments. The level of significance was set at $P \leq 0.05$. CD (Critical difference) was used to check the significance of the results (Sheoran and Pannu, 1999).

3. RESULTS AND DISCUSSION

3.1 Organoleptic Acceptability

Laddo developed from oilseed meals were found organoleptically acceptable by panel of judges. *Laddo* prepared without oilseed meal served as control. The mean acceptability scores of control *laddo* were 8.87, 8.70, 8.70, 8.50, 8.63, and 8.68 for color, appearance, aroma, texture, taste, and overall acceptability, respectively (Table 2).

3.2 Sesame Meal *Laddo*

The mean acceptability scores for color, appearance, aroma, texture and taste were ranged from 8.33 to 8.38, 8.10 to 8.40, 8.09 to 8.26, 7.91 to 8.00, and 8.18 to 8.30, respectively. Results elucidates that overall acceptability scores for T1, T2 and T3 *laddo* were 8.27, 8.21, and 8.12, respectively. The taste, texture, appearance, and overall acceptability of all sesame meal *laddo* were ranged from 'liked moderately' to 'liked very much' category and control *laddo* sensory attributes lied in 'liked very much' category (Table 2).

3.3 Flaxseed Meal *Laddo*

The mean acceptability scores for developed *laddo* ranged between 8.34 to 8.48, 8.20 to 8.60, 8.12 to 8.40, 7.88 to 8.04 and 8.04 to 8.23 for color, appearance, aroma, texture and taste, respectively. Overall acceptability scores for T1 (12%), T2 (18%) and T3 (24%) *laddo* was 8.41, 8.18 and 8.12, respectively (Table 2).

Table 2. Organoleptic acceptability of *laddo* incorporating sesame, flaxseed, groundnut, and composite meal (Mean scores)

Types of <i>laddo</i>		Color	Appearance	Aroma	Texture	Taste	Overall acceptability
Control	Control	8.87±0.06	8.70±0.09	8.70±0.09	8.50±0.08	8.63±0.06	8.68±0.14
Sesame meal	T1	8.38±0.09	8.40±0.1	8.26±0.15	8.00±0.01	8.30±0.15	8.27±0.1
	T2	8.38±0.19	8.30±0.13	8.16±0.21	7.98±0.05	8.24±0.21	8.21±0.09
	T3	8.33±0.07	8.10±0.17	8.09±0.16	7.91±0.07	8.18±0.06	8.12±0.12
Flaxseed meal	T1	8.48±0.11	8.60±0.15	8.40±0.23	8.04±0.06	8.23±0.11	8.41±0.11
	T2	8.40±0.13	8.20±0.08	8.19±0.19	7.92±0.08	8.21±0.15	8.18±0.06
	T3	8.34±0.13	8.20±0.2	8.12±0.15	7.88±0.08	8.04±0.13	8.12±0.05
Groundnut meal	T1	8.38±0.07	8.30±0.1	8.68±0.1	7.98±0.01	8.14±0.02	8.30±0.16
	T2	8.36±0.03	8.10±0.12	8.17±0.04	7.91±0.06	7.91±0.02	8.09±0.05
	T3	8.31±0.16	8.10±0.03	8.11±0.03	7.88±0.03	7.90±0.07	8.06±0.09
Composite meal	C1	8.39±0.11	8.80±0.13	8.70±0.15	8.42±0.05	8.80±0.12	8.62±0.19
	C2	8.34±0.13	8.80±0.15	8.60±0.06	8.27±0.07	8.66±0.02	8.53±0.2
	C3	8.30±0.1	8.70±0.15	8.45±0.06	8.00±0.1	8.41±0.04	8.37±0.19
C.D.≤0.05		N/A	0.37	0.41	0.18	0.31	N/A

Values are mean of ten panelists

Control- without any oilseed meal incorporation (T1, T2 and T3 are 12%, 18%, and 24%)

3.4 Groundnut Meal Laddo

Three types of groundnut meal based *laddo* were prepared. Mean acceptability scores for color, appearance, aroma, texture, taste, and overall acceptability of T1 *laddo* were 8.38, 8.30, 8.68, 7.98, 8.14, and 8.30, respectively. Mean acceptability scores for color, appearance, aroma, texture, taste, and overall acceptability of T2 *laddo* were 8.36, 8.10, 8.17, 7.91, 7.91, and 8.09, respectively. Mean acceptability scores for color, appearance, aroma, texture, taste, and overall acceptability of T3 *laddo* were 8.31, 8.10, 8.11, 7.88, 7.90, and 8.06, respectively (Table 2).

3.5 Composite Meal Laddo

All the three oilseed meals mixed in equal proportion and used as composite meal which was incorporated at 12, 18 and 24% level in *laddo*. Result delineates that mean acceptability scores for C1 (12%) *laddo* were 8.39, 8.80, 8.70, 8.42, 8.80, and 8.62 for color, appearance, aroma, texture, taste, and overall acceptability, respectively. Mean acceptability scores for color, appearance, aroma, texture, taste, and overall acceptability of C2 (18%) *laddo* were 8.34, 8.80, 8.60, 8.27, 8.66, and 8.53, respectively. Mean acceptability scores for C3 (24%) *laddo* were 8.30, 8.70, 8.45, 8.00, 8.41, and 8.37 for color, appearance, aroma, texture, taste, and overall acceptability, respectively whereas mean acceptability scores of control *laddo* were 8.87, 8.70, 8.70, 8.50, 8.63, and 8.68 for color, appearance, aroma, texture, taste, and overall acceptability, respectively (Table 2).

All types of supplemented *laddo* were organoleptically acceptable and overall acceptability of each *laddo* lied in 'liked very much' so, the highest level of incorporation level were selected and studied for nutritional analysis.

3.6 Nutritional Evaluation

The moisture content of all *laddo*'s lied between 0.05 to 0.96%. The crude protein was observed

to be 20.03% in groundnut meal *laddo*, 18.34% in composite meal *laddo*, 16.58% in sesame meal *laddo*, 13.35% in flaxseed meal *laddo* and 9.16% in control *laddo*. However, control *laddo* fat content was 23.54% which was significantly ($P \leq 0.05$) low. The values of ash content of oilseed incorporated *laddo* varied between 2.44% to 5.70%. The amount of crude fiber was found 4.28% in flaxseed meal *laddo*, 4.19% in groundnut meal *laddo* and 2.65% in control *laddo* whereas in composite meal *laddo* crude fiber content was 5.14% and in sesame meal *laddo* crude fiber content was 4.85%. Significantly ($P \leq 0.05$) low values were observed for crude protein, crude fat, ash and crude fiber of control *laddo* as compared to oilseed meal based *laddo* (Table 3).

Comparison of protein and moisture of all the selected oilseed meal *laddo* indicated that it was significantly ($P \leq 0.05$) high in groundnut meal. Sesame and flaxseed had almost similar protein. Comparison of fat content of all the oilseed meal *laddo* indicated that crude fat content of flaxseed meal *laddo* was significantly high (35.99%) as compared to sesame meal, groundnut meal and composite meal *laddo*. High fat content in flaxseed meal *laddo* was due to fact that oil from flaxseed was extracted with cold pressed method in which the oil retention is more in meal. Control *laddo* had lowest fat i.e. 23.45%. Significant ($P \leq 0.05$) difference observed in fat content among all oil seed meal *laddo* may be due to fat content of oil seed and method of oil extraction used. The comparison of ash content among sesame, flaxseed, groundnut and composite meal *laddo* revealed that ash content was significantly ($P \leq 0.05$) high in sesame meal *laddo* and lowest in control *laddo*. A variation was observed in crude fiber content among all oil seed meal *laddo* evaluated. Maximum amount of crude fiber was found in composite meal *laddo* and lowest found in control *laddo* whereas flaxseed meal and groundnut meal *laddo* had almost similar amount of crude fiber.

Table 3. Proximate composition of oilseed meal based *laddo* (% , on dry matter basis)

Types of <i>Laddo</i>	Moisture	Crude Protein	Crude Fat	Ash	Crude Fiber
Control	0.96±0	9.16±0.09	23.54±0.31	2.44±0.01	2.65±0
Sesame meal	0.14±0	16.58±0.38	27.8±0.52	5.7±0.02	4.85±0
Flaxseed meal	0.05±0	13.35±0.02	35.99±0.65	3.97±0.01	4.28±0.08
Groundnut meal	0.41±0	20.03±0.45	25.52±0.07	4.66±0.06	4.19±0.04
Composite meal	0.08±0	18.34±0.27	30.6±0.43	4.19±0.02	5.14±0.07
C.D. ($P \leq 0.05$)	0.01	0.94	1.42	0.1	0.17

Values are mean±SE of three independent determinations
* Moisture calculated on wet matter basis; * SE=Standard error

Table 4. Nutritional composition of oilseed meals (Percent)

Sesame meal					
Sr. no.	Protein	Fat	Fibre	Ash	Reference
1.	35.16	5.52	23.52	9.98	Hira et al. (2002)
2.	41.04	8.05	6.1	11.13	Nadeem et al. (2005)
3.	44.42	13.11	8.75	14.15	Babiker (2012)
4.	51.55	6.24	9.59	11.34	Son et al. (2017)
5.	44.9	11.3	7.3	11.9	Heuze et al. (2017)
Flaxseed meal					
1.	41.12	5.63	8.76	5.38	Bell et al. (1993)
2.	32.9	9.1	13.2	6.6	De Boever et al. (1994)
3.	34.56	6.3	10.01	6.5	Masoero et al. (1994)
4.	38.00-47.30	12.8-26.1	-	3.7-5.1	Ogunronbi et al. (2011)
5.	38.4	12.8	7.2	5.1	Kyntaja et al. (2014)
6.	34.1	10.2	10.5	6.3	Heuze et al. (2017)
Groundnut meal					
1.	45.1	9.9	4	7.9	Awoniyi et al. (2003)
2.	38.59	8.55	7.52	6.13	Chandrasekharaiah et al. (2004)
3.	52.45	10.56	4.8	5.5	Adewolu et al. (2010)
4.	42.87	7.75	--	5.51	Gupta et al. (2011)
5.	53.44	7.47	8.55	5.27	Babiker (2012)
6.	49.1	9.8	7	5.8	Heuze et al. (2018)

* The mean values of proximate composition of sesame meal, flaxseed meal, and groundnut meal obtained by mechanical extraction as reported by various researchers in different studies

The experiment was planned due to lack of literature available for traditional food products based on oilseed meals and the results obtained for organoleptic acceptability were highly acceptable. However, there was no data available for the developed products from oilseed meal, although data on the proximate composition of oilseed meals were available (Table 4).

4. CONCLUSION

One serving (25 gm) of *laddo* can meet the 27% requirement of protein for pre-schoolers as per RDA (recommended dietary allowance). It can be summarized from the present study that the *laddo* made using oilseed meals is nutrient-dense which contains protein, fat, ash, crude fiber content in ample amounts. The study has shown a potential use of inexpensive and underutilized oilseed meals in the preparation of various food products for human consumption. The developed value-added *laddo* could be efficiently used as a means of improving the nutritional status of the community by popularizing and making them reach the masses to overcome malnutrition. The *laddo* if promoted will also help in the utilization of oilseed meals which are generally discarded. Sesame meal,

flaxseed meal, groundnut meal and composite meal of different meals can be used to supplement foods consumed as a part of daily diet.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The Author(s) hereby declares that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

Authors would like to acknowledge the financial help received from CSHAU, Hisar as PG scholarship throughout the research. We wish to acknowledge laboratory of experiential learning and Food Analysis, Department of Foods and Nutrition, CCS, Haryana Agricultural University, Hisar.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Adewolu, M.A., Ikenweibe, N.B., and Mulero, S.M. (2010). Evaluation of an animal protein mixture as a replacement for fish meal in practical diets for fingerlings of *Clarias gariepinus* (Burchell, 1822). *Israeli Journal of Aquaculture Bamidgeh*, 62(4), 237–244.
- AOAC. Official Methods of Analysis. Association of Official Analytical Chemist. Washington, D.C; 2010.
- Awoniyi, T.A.M., Aletor, V.A., and Aina, J.M. (2003). Performance of broiler-chickens fed on maggot meal in place of fishmeal. *International Journal of Poultry Science*.
- Babiker, M.S. (2012). Chemical composition of some non-conventional and local feeder sources for Poultry in Sudan. *International Journal of Poultry Science*, 11(4), 283–287.
- Bell, J.M., and Keith, M.O. (1993). Nutritional evaluation of linseed meals from flax with yellow or brown hulls, using mice and pigs. *Animal Feed Science and Technology*, 43(1–2), 1–18.
- Chandrasekharaiah, M., Sampath, K.T., Praveen, U.S., Umalatha, (2004). Chemical composition and *in vitro* digestibility of certain commonly used feed stuffs in ruminant rations. *Indian J. Dairy Sci.*, (57), 114-117.
- De Boever, J.L., Cottyn, B.G., Vanacker, J.M., and Boucque, C.V. (1994). An improved enzymatic method by adding gamma-nase to determine digestibility and predict energy value of compound feeds and raw materials for cattle. *Animal Feed Science and Technology*, 47(1–2), 1–18.
- Gupta, A., Singh, S., Kundu, S.S., and Jha, N. (2011). Evaluation of tropical feedstuffs for carbohydrate and protein fractions by CNCV system. *Indian Journal of Animal Sciences*, 81(11), 1154–1160.
- Heuze, V., Tran, G., Bastianelli, and D., Lebas, F., (2017). Sesame (*Sesamum indicum*) seeds and oil meal. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO.
- Heuze, V., Tran, G., Noziere, P., Lessire, M., Lebas, F. 2018. *Linseed. Feedipedia, a programme by INRAE, CIRAD, AFZ, and*
- FAO.
meal<https://www.feedipedia.org/node/735>
- Hira, A.K., Ali, M.Y., Chakrabort, M., Islam, M.A., and Zaman, M.R. (2002). Use of Waterhyacinth Leaves (*Eichhorina crassipes*) Replacing Dhal Grass (*Hymenachne pseudo interrupta*) in the Diet of Goat. *Pakistan Journal of Biological Sciences*, 5(2), 218–220.
- Karnika, Kawatra, A., Verma, J., Rani, S., Deepankar, & Kumar, S. (2022). Nutrition evaluation of oilseed meals. *Journal of Agriculture Research and Technology*, 47(4, Special Issue), 45–56.
- Kyntaja, S., Partanen, K., Siljander-Rasi, H., and Jalava, T. (2014). *Tables of composition and nutritional values of organically produced feed materials for pigs and poultry*. MTT Report 164.
- Masoero, F., Fiorentini, L., Rossi, F., and Piva, A. (1994). Determination of nitrogen intestinal digestibility in ruminants. *Animal Feed Science and Technology*, 48(3–4), 253–263.
- Nadeem, M.A., Gilani, A.H., Khan, A.G., and Mahr-Un-Nisa. (2005). True Metabolizable Energy values of poultry feedstuffs in Pakistan. *International Journal of Agriculture and Biology*, 7(6), 990–994.
- Ogunronbi, O., Jooste, P.J., Abu, J.O., and Van der Merwe, B. (2011). Chemical composition, storage stability and effect of cold-pressed flaxseed oil cake inclusion on bread quality. *Journal of Food Processing and Preservation*, 35(1), 64–79.
- Park, C.S., Helmbrecht, A., Htoo, J.K., and Adeola, O. 2017. Comparison of amino acid digestibility in full-fat soybean, two soybean meals, and peanut flour between broiler chickens and growing pigs. *Journal of Animal Science*, 95(7), 3110–3119.
- Sheoran, O.P. and Pannu, R.S. 1999. Statistical Package for agricultural workers. "O. P. Stat" College of Agriculture, Kaul, CCS Haryana Agricultural University, Hisar. India
- Son, A.R., Park, C.S., and Kim, B.G. (2017). Determination and prediction of digestible and metabolizable energy concentrations in byproduct feed ingredients fed to growing pigs. *Asian-Australasian Journal of Animal Sciences*, 30(4), 546–553.

The Bloomsbury Handbook of Indian Cuisine.
Bloomsbury Publishing. 2023. pp. 269–
270. ISBN 978-1-350-12865-1. Archived

from the original on 2023-03-14. Retrieved
2023-03-14.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/129108>